

# Coupled Electron-Ion Monte Carlo Study of Hydrogen

Kris Delaney and David Ceperley

*Department of Physics, University of Illinois at Urbana-Champaign  
Urbana, IL, USA*

Carlo Pierleoni

*Dipartimento di Fisica, Università del L'Aquila  
L'Aquila, Italy*

We present details of the Coupled Electron-Ion Monte Carlo method (CEIMC)<sup>1,2</sup> applied to the problem of the equation of state of pure hydrogen. The aim is to develop a method that can predict state information outside the range of temperatures and pressures that are accessible with other existing methods, such as PIMC.

The CEIMC method centers on exploring the configuration space of the hydrogen nuclei (classical or quantum path integrals) using a modified Metropolis algorithm, with configurational energy differences computed from Born-Oppenheimer energies. Energy differences are computed with VMC or Reptation quantum Monte Carlo, both of which supply unbiased estimates of energy differences, the latter within a projector framework.

Using a wavefunction generated through a band-structure calculation, we study the so-called plasma phase transition, the conditions under which a fluid of pure molecular hydrogen dissociates.

- [1] D. Ceperley, M. Dewing and C. Pierleoni, in *Bridging Time Scales: Molecular Simulations for the Next Decade*, eds. P. Nielaba, M. Mareschal and G. Ciccotti, Springer-Verlag, pgs. 473-500 (2002).
- [2] C. Pierleoni, D. M. Ceperley and M. Holzmann, *Phys. Rev. Lett.* **93**, 146402 (2004)