

Liquid-liquid transition in high pressure hydrogen by Quantum Monte Carlo methods

Carlo Pierleoni^{*1,2}

¹Department of Physical and Chemical Sciences, University of L'Aquila, Italy

²Maison de la Simulation, CEA-Saclay, France

We studied the transformation from molecular-insulating to monatomic-metallic fluid hydrogen upon increasing pressure, by first principle simulations based on both Density Functional Theory (DFT) and Quantum Monte Carlo (QMC) methods [1,2]. Below a critical temperature $T_c \in [1500K, 3000K]$, the transition is first order with a discontinuity in the specific volume, a sudden dissociation of the molecules and a discontinuous change in electronic properties like the momentum distribution and the electrical conductivity [3,4,5]. Above the critical point the transformation into the metallic dissociated state is continuous. Optical response of the system is obtained within the Kubo-Greenwood framework of DFT. Agreement with recent experimental results at NIF for reflectivity and absorption is observed. Our analysis suggest a coherent picture for apparently discordant experiments from different methods.

- [1] C. Pierleoni, M.A. Morales, G. Rillo, M. Holzmann and D.M. Ceperley, "Liquid-liquid phase transition in hydrogen by Coupled Electron-Ion Monte Carlo Simulations", PNAS 113, 4953-4957(2016).
- [2] M.A. Morales, C. Pierleoni, E. Schwegler and D.M. Ceperley, "Evidence for a first-order liquid-liquid transition in high-pressure hydrogen from ab initio simulations", PNAS 107, 12799 (2010).
- [3] C. Pierleoni, M. Holzmann and D.M. Ceperley, "Local structure in dense hydrogen at the liquid-liquid phase transition by coupled electron-ion Monte Carlo", Contrib. Plasma Phys. 58, 99-106 (2018).
- [4] C. Pierleoni, G. Rillo, D.M. Ceperley and M. Holzmann, "Electron localization properties in high pressure hydrogen at the liquid-liquid phase transition by Coupled Electron-Ion Monte Carlo", J. Phys.: Conf. Ser. 1136, 012005 (2018).
- [3] V. Gorelov, C. Pierleoni and D.M. Ceperley, "Benchmarking vdW-DF first-principles predictions against Coupled Electron-Ion Monte Carlo for high-pressure liquid hydrogen", Contributions to Plasma Physics (2019) ;e201800185. <https://doi.org/10.1002/ctpp.201800185>.
- [5] G. Rillo, M.A. Morales, D.M. Ceperley and C. Pierleoni, "Optical properties of high pressure liquid hydrogen across molecular dissociation", PNAS (2019), in print, arXiv:1810.08131v1.
- [6] P. Celliers et al, "Insulator-metal transition in dense fluid deuterium", Science 361, 677 (2018).

*carlo.pierleoni@aquila.infn.it,carlo.pierleoni@cea.fr